

## Effect of an avocado oil-rich diet over an angiotensin II-induced blood pressure response

M.J. Salazar<sup>a</sup>, M. El Hafidi<sup>b</sup>, G. Pastelin<sup>a</sup>, M.C. Ramírez-Ortega<sup>a</sup>,  
M.A. Sánchez-Mendoza<sup>a,\*</sup>

<sup>a</sup> Instituto Nacional de Cardiología “Ignacio Chávez”, Department of Pharmacology, Juan Badiano No. 1, Col. Sección XVI, Deleg. Tlalpan, Distrito Federal, C.P. 14080, México

<sup>b</sup> Instituto Nacional de Cardiología “Ignacio Chávez”, Department of Biochemistry, Juan Badiano No. 1, Col. Sección XVI, Deleg. Tlalpan, Distrito Federal, C.P. 14080, México

Received 1 November 2003; received in revised form 2 January 2005; accepted 19 January 2005

### Abstract

We studied the effect of an avocado oil-rich diet on (1) the blood pressure response to angiotensin II (AngII) and (2) the fatty acid composition of cardiac and renal membranes on male Wistar rats. The avocado oil-rich diet induced a slightly higher AngII-induced blood pressure response in the rats as compared to the control rats. In cardiac microsomes, avocado oil induced an increase in oleic acid content ( $13.18 \pm 0.33\%$  versus  $15.46 \pm 0.59\%$ ), while in renal microsomes, the oil decreased  $\alpha$ -linolenic acid content ( $0.34 \pm 0.02\%$  versus  $0.16 \pm 0.12\%$ ), but increased the arachidonic acid proportion ( $24.02 \pm 0.54\%$  versus  $26.25 \pm 0.54\%$ ), compared to control. In conclusion, avocado oil-rich diet modifies the fatty acid content in cardiac and renal membranes in a tissue-specific manner. The rise in renal arachidonic acid suggests that diet content can be a key factor in vascular responses.

© 2005 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Avocado oil; Blood pressure; Membrane fatty acid composition; Arachidonic acid; Angiotensin II

### 1. Introduction

*Persea americana* Miller (Lauraceae) or avocado, native to Mexico, is valued due to its nutritional and therapeutic qualities. Avocado fruit and leaves have been used in Mexican folk medicine to treat a wide variety of diseases. Francisco Hernández reported as early as the XVI century, that oil obtained from pressing the seed was useful in the treatment of rashes and scars, had an astringent effect, and could also be used to treat dysentery (Argueta-Villamar et al., 1994). Hot water infusion from the leaves can be taken as an emmenagogue, diuretic, to treat coughs and colds, and diarrhea. An important use of avocado leaves is to treat hypertension, which is not used exclusively by Mexican populations, but

people in countries like Brazil, Indonesia, Jamaica, Nigeria and Panama also use it for the same therapeutic benefits (Ross, 1999; Adeboye et al., 1999).

Recent studies carried out among Mexican populations that consume avocados have shown that the avocado decreases serum total cholesterol, LDL-cholesterol and triglycerides, and increases HDL-cholesterol levels compared to the control diet (Alvizouri-Munoz et al., 1992; Lopez-Ledesma et al., 1996).

The consumption of oil from different sources exerts different effects over the lipid composition of the cellular membranes and their function. Therefore, we studied the effect of a diet rich in avocado oil and a control diet, on the blood pressure response to angiotensin II (AngII). In addition, we evaluated the effect of an avocado oil diet on fatty acid composition of cardiac and renal membranes in order to correlate biochemical changes and physiological responses.

\* Corresponding author. Tel.: +52 55 5573 2911x1317; fax: +52 55 5573 0926.

E-mail address: [masanchez@gmail.com](mailto:masanchez@gmail.com) (M.A. Sánchez-Mendoza).

## 2. Materials and methods

### 2.1. Experimental animals

Male Wistar rats (230–250 g), bred and raised in our facilities, were placed into metabolic cages 3 days prior to the beginning of the protocol, offered tap water and lab chow *ad libitum*, and maintained on a 12-h light:12-h dark cycle in a temperature-controlled room. Animals were randomly divided into two experimental groups of five rats each. The control group received lab chow, while the treated group received a 10% (w/w) avocado oil-rich diet for a 2-week period. At the end of the treatment, the rats were anaesthetized with sodium pentobarbital (50 mg/kg, *i.p.*) and were either prepared for blood pressure measurement, or the heart and kidneys were removed. All procedures were conducted in accordance with Institutional ethical guidelines.

### 2.2. Diet preparation

The avocado-enriched diet was prepared in bulk in our laboratory by mixing ground lab chow with avocado oil obtained from fresh fruit (10%, w/w). Both the control diet (ground lab chow) and avocado oil-rich diet were partitioned into daily rations packaged in plastic bags, and flushed with nitrogen to minimize oxidation and stored at 4 °C.

### 2.3. Fatty acid methyl esters analysis

Cardiac and renal microsomes were obtained as described by Garg and co-workers (1988) in a homogenizing buffer containing: 250 mM sucrose, 0.1 mM ethylenediamine tetraacetic acid (EDTA), 62 mM potassium phosphate, 150 mM potassium chloride, 5 mM magnesium chloride, and 1 mM dithiothreitol (DTT), at pH 7.4. The microsomes, containing butylated hydroxy toluene (BHT, 0.02%), were stored at –70 °C until processed. The lipids were extracted as described by Folch et al. (1957). The lipid extracts were trans-esterified to their fatty acid methyl esters as described by Christie (1989), separated and identified by gas–liquid chromatography in a Carlo Erba Fractovap 2300 chromatograph, fitted with a 25 m × 0.25 mm i.d. fused-silica capillary column, coated with CP-Sil 88 (film thickness, 0.25 µm). The analysis was carried out at an isotherm temperature of 195 °C, using helium gas as a carrier at a flow rate of 1 ml/min.

### 2.4. Angiotensin II-induced increase of blood pressure

At the end of the 2-week treatment, both control rats and rats fed with avocado oil-rich diet were anaesthetized and we performed intra-artery measurement of the blood pressure as described by Adeboye et al. (1999). AngII (100, 300, and 1000 ng/kg, *i.v.*) was administered and the basal- and AngII-induced blood pressure changes were measured with a Blood Pressure monitor (BP Monitor, WPI, USA).

### 2.5. Statistical analysis

Data were expressed as mean ± S.E.M. Statistical evaluation of the data was performed using Student's *t*-test for unpaired comparisons;  $p < 0.05$  was considered statistically significant.

## 3. Results

### 3.1. Effect on body weight

Two weeks of an avocado oil-rich diet had no significant influence on the rat's body weight. Body weight of control rats was  $247.5 \pm 1.4$  g, while in the avocado oil-rich diet was  $249.1 \pm 1.7$  g.

### 3.2. Effect on AngII-induced change in blood pressure

Basal systolic blood pressure for control and avocado oil-rich diet rats was:  $95 \pm 3.1$  and  $97 \pm 2.6$  mmHg, respectively. The administration of AngII (100, 300, and 1000 ng/kg, *i.v.*) induced an increase in blood pressure in both control rats and rats fed with avocado oil-rich diet; however, this increment was slightly higher in the rats fed with avocado oil-rich diet, as compared to the control rats (Fig. 1).

### 3.3. Effect on fatty acid composition

Avocado oil-rich diet induced an increase in oleic acid proportion in cardiac microsomes ( $15.46 \pm 0.5\%$  of total) compared to those from control rats ( $13.18 \pm 0.3\%$ ,  $p < 0.05$ ). In renal microsomes, avocado oil-rich diet elicited a decrease in  $\alpha$ -linolenic acid ( $0.34 \pm 0.02$  and  $0.16 \pm 0.12\%$  of total, for control and avocado oil-treated rats, respectively,  $p < 0.05$ ); as well as an increase in arachidonic acid proportion: for control,




Fig. 1. Effect of angiotensin II-stimulation on blood pressure response in rats fed a control and avocado oil-rich diet. AngII (100, 300, and 1000 ng/kg, *i.v.*) was administered to rats fed control diet (■), and avocado oil-rich diet (10%, w/w, □) for 2 weeks. The bars represent the mean ± S.E. of five different experiments.

Download English Version:

<https://daneshyari.com/en/article/9009810>

Download Persian Version:

<https://daneshyari.com/article/9009810>

[Daneshyari.com](https://daneshyari.com)