



## Relationships between depression, gender, and unhealthy weight loss practices among overweight or obese college students<sup>☆</sup>



E.P. Davila<sup>a,b,\*</sup>, J.K. Kolodziejczyk<sup>a,b,c</sup>, G.J. Norman<sup>a,b</sup>, K. Calfas<sup>b</sup>, J.S. Huang<sup>a,b,d,e</sup>, C.L. Rock<sup>b</sup>, W. Griswold<sup>f</sup>, J.H. Fowler<sup>a,g</sup>, S.J. Marshall<sup>a,b</sup>, A. Gupta<sup>a</sup>, K. Patrick<sup>a,b</sup>

<sup>a</sup> Center for Wireless & Population Health Systems (CWPHS), Qualcomm Institute/Calit2, University of California, San Diego, 9500 Gilman Drive, Atkinson Hall, Dept 0811, La Jolla, CA 92093, USA

<sup>b</sup> Department of Family & Preventive Medicine, University of California, San Diego, 9500 Gilman Drive, Dept 0628, La Jolla, CA 92093, USA

<sup>c</sup> Graduate School of Public Health, San Diego State University, 5500 Campanile Drive, Hardy Tower 119, San Diego, CA 92182, USA

<sup>d</sup> Rady Children's Hospital, 3020 Children's Way, San Diego, CA 92123, USA

<sup>e</sup> Department of Pediatrics, University of California, San Diego, 3020 Children's Way, MC 5109, La Jolla, CA 92093, USA

<sup>f</sup> Department of Computer Science & Engineering, University of California, San Diego, 9500 Gilman Drive, Dept 0404, La Jolla, CA 92093, USA

<sup>g</sup> Medical Genetics Division & Political Science Department, University of California, San Diego, 9500 Gilman Drive, Dept 0521, La Jolla, CA 92093, USA

### ARTICLE INFO

#### Article history:

Received 18 September 2013

Received in revised form 10 January 2014

Accepted 11 March 2014

Available online 27 March 2014

#### Keywords:

Weight loss

Men

Women

Young adult

Depression

Eating disorder

### ABSTRACT

**Introduction:** Unhealthy weight loss practices are common among female college students. It is unknown if these practices are also most common among women in the subset of overweight or obese college students or if these practices are related to depression. We examined the relationship between gender, depression, and unhealthy weight loss practices among overweight or obese college students.

**Methods:** Students (body mass index between 25.0 and 34.9 kg/m<sup>2</sup>) from three Southern California universities ( $M_{age} = 22$  years,  $SD = 4$ ; 70% women) were recruited from May 2011 to May 2012 for participation in a weight loss clinical trial ( $N = 404$ ). Logistic regressions were performed with baseline data to assess the cross-sectional relationship between self-reported unhealthy weight loss practices and gender and depression as measured by the Center for Epidemiologic Studies Depression short form.

**Results:** Twenty-nine percent of participants reported engaging in at least one unhealthy weight loss behavior (e.g., fasting, purging) over the last 30 days, with no differences by gender. Self-report of at least one unhealthy weight loss behavior was associated with report of symptoms of depression ( $e^B = 1.14$  [confidence interval, CI: 1.08–1.20]), adjusting for potential confounders. Interactions between gender and depression were not significant ( $e^B = 1.04$  [CI: 0.93–1.16]).

**Conclusion:** Among an overweight or obese sample of college students, unhealthy weight loss practices were equally common in both genders, and students with depressive symptomatology were at greatest risk. Obesity interventions targeting overweight or obese college students should educate both men and women about the dangers of unhealthy weight loss practices. In addition, screening for depression can help identify students who would benefit from additional supportive and coping strategies and resources.

© 2014 Elsevier Ltd. All rights reserved.

### 1. Introduction

Research suggests that unhealthy weight loss practices (e.g., fasting, vomiting, taking diet pills, laxatives, or diuretics, smoking cigarettes as food substitute) are common among college students, particularly college women (Bish et al., 2005; Harring, Montgomery, & Hardin, 2011; Striegel-Moore et al., 2009). In the 2008 National College Health

Assessment (NCHA) survey of 90,484 United States (U.S.) college students, women were more likely than men to report dieting to lose weight (42.1% vs. 24.1%), vomiting or using laxatives (3.3% vs. 0.6%), and taking diet pills (3.8% vs. 1.7%); these estimates included normal to obese students (Harring et al., 2011). Depression may be one possible reason why college students, particularly women, engage in unhealthy weight loss practices. Harring et al. (2011) found that among overweight college students with an accurate weight perception, women were more likely than males to be depressed (Harring et al., 2011). In a study of 2822 U.S. college students, a greater proportion of women than men screened positive for both an eating disorder and depression (Eisenberg, Nicklett, Roeder, & Kirz, 2011). In fact, eating disorders often are associated with depression among college women (Eisenberg et al., 2011; Gutzwiller, Oliver, & Katz, 2003; Meno, Hannum, Espelage, & Low,

*Abbreviations:* CESD-10, Center for Epidemiologic Studies Depression short form; SD, standard deviation.

<sup>☆</sup> Clinical trial registration number: #NCT01200459.

\* Corresponding author at: Department of Family & Preventive Medicine, University of California, San Diego, 9500 Gilman Drive, Dept. 0811, La Jolla, CA 92093-0811, USA. Tel.: +1 858 534 9550; fax: +1 858 534 9404.

E-mail address: evelyn.p.davila@gmail.com (E.P. Davila).

2008). This may be because college women are most vulnerable to social norms for thinness, with many of these women suffering from low self-esteem (Elgin & Pritchard, 2006; Harring et al., 2011; Klesges, Mizes, & Klesges, 1987). Because eating disorder symptoms also have been associated with depression among college men (Olivardia, Pope, Mangweth, & Hudson, 1995), their possible depression is associated with unhealthy weight loss practices regardless of gender. Nevertheless, because more college women than men have been found to diet to lose weight at lower BMIs (Bish et al., 2005; Harring et al., 2011), it is possible that among a sample of overweight and obese college students, gender differences in engagement in unhealthy weight loss practices will not be evident.

The purpose of this analysis was to assess the association between gender, depression, and unhealthy weight loss practices in a sample of overweight or obese college students from Southern California. We hypothesized that 1) men will be as likely as women to engage in at least one unhealthy weight loss practice, 2) engagement in at least one unhealthy weight loss practice will be associated with greater symptoms of depression, 3) the strength of association between depression and engagement in at least one unhealthy weight loss practice will be greater among women than men.

## 2. Methods

### 2.1. Participants

The current analyses used baseline data from a randomized controlled trial to assess a theory-based weight loss program delivered via social and mobile technologies, project SMART (SMART: Social Mobile Approaches to Reduce weight) (Patrick et al., 2013). College students ( $N = 404$ ) between the ages of 18 and 35 years with a BMI between 25.0 and 34.9 kg/m were recruited from May 2011 to May 2012. Participants were students enrolled at San Diego State University, California State University San Marcos, or University of California San Diego. Institutional review boards from these universities approved study protocols. Individuals were excluded if they: 1) met American Diabetes Association criteria for diabetes; 2) screened positive for an eating disorder or a cardiovascular condition; 3) were taking weight-altering medications; 4) were pregnant or intending to get pregnant over the study period; and 5) were enrolled in or planned to enroll in another weight loss program. More information about project SMART can be found elsewhere (Patrick et al., 2013).

### 2.2. Measures

During the baseline visit, physiologic measurements were taken, and participants completed surveys with questions about demographics, unhealthy weight loss practices, and depression symptoms.

Body weight was measured to the nearest 0.1 kg using a calibrated digital scale. Subjects were asked to wear lightweight clothes (e.g., exercise clothes). Height (without shoes) was measured to the nearest 0.1 cm using a stadiometer. BMI was calculated from height and weight ( $\text{kg}/\text{m}^2$ ). Those with a BMI between 25.0 and 29.9  $\text{kg}/\text{m}^2$  were classified as overweight while those with  $\text{BMI} \geq 30 \text{ kg}/\text{m}^2$  were classified as obese (Centers for Disease Control and Prevention, 2011). The Seca703, a combined digital scale and stadiometer, was used for body weight and height measurements.

Demographic characteristics included the following: age (continuous), race (1: White, 2: Black or African American, 3: Asian, 4: Pacific Islander, 5: American Indian, 6: Other), Hispanic ethnicity (1: Yes, 2: No), college graduate (1: Yes, 2: No), and relationship status (1: Single, 2: Engaged, married or in a committed relationship, 3: Separated, divorced, or widowed).

Engagement in unhealthy weight loss practices was based on the question: "Over the last 30 days, have you done any of the following things in order to lose weight or to keep from gaining weight?" The

list included: fasting, eating very little food, taking diet pills, self-vomiting, using laxatives, using diuretics, skipping meals, or smoking cigarettes. A score was created using the sum of the 'yes' responses. A dichotomous variable also was created and defined as the engagement in at least one unhealthy weight loss practice. These items and dichotomization have been used by others and show high test–retest agreement (Neumark-Sztainer, Wall, Larson, Eisenberg, & Loth, 2011).

Depression symptomatology (e.g., depressed mood, feelings of guilt, worthlessness and helplessness, psychomotor retardation, loss of appetite, or sleep difficulties) was based on self-report and measured with the Center for Epidemiologic Studies Depression (CESD) short form (CESD-10) (Andresen, Malmgren, Carter, & Patrick, 1994). This short form is a 10-item subset of the 20-item CESD scale and has a 30-point categorical response scale (range 0–30). Higher scores represent greater depressive symptoms, and a score of  $\geq 10$  is indicative of depression; this threshold was used to create a dichotomous variable for depression. The widely used CESD has shown good reliability and validity, including among college students (Carleton et al., 2013; Radloff, 1991). The short form has shown good reliability and validity as well (Andresen et al., 1994), but its psychometric properties among college students are unknown.

### 2.3. Statistical analyses

Chi-square and t-tests were used to assess gender differences in demographics, BMI, CESD score, and engagement in unhealthy weight loss practices. The unadjusted association between gender and CESD score (continuous variable) and engagement in at least one unhealthy weight loss practice was assessed with a univariable logistic regression, with engagement in at least one unhealthy weight loss practice as the outcome variable. The unadjusted association between CESD score and gender was assessed using univariable linear regression and CESD score as the outcome variable. The CESD score  $\times$  gender interaction was tested. Associations and interactions with BMI were also tested given findings by others (Harring et al., 2011; Hoerr, Bokram, Lugo, Bivins, & Keast, 2002).

The multivariable hierarchical logistic regression model included, in addition to gender and CESD score, demographics and BMI, as these may be potential confounders (Hoerr et al., 2002; Madanat, Hawks, & Novilla, 2006). To increase statistical power of the regression analysis, race categories of Pacific Islander, American Indian, and others were collapsed; the relationship status categories of single and separated, divorced or widowed were collapsed; and the continuous variables for CESD score and BMI were used (instead of dichotomous variables). Variables were considered significant at  $p < 0.05$ . The Hosmer–Lemeshow Goodness-of-Fit statistic was used to assess model fit. An alpha level of 0.05 was used for all statistical tests.

## 3. Results

Table 1 shows participant demographics and health status by sex. Many participants were women, single, White, undergraduates, overweight, and/or without depressive symptoms; 11% of women and 12% of men were overweight and met criteria for depression (i.e., CESD score  $\geq 10$ ). There were no statistically significant differences between men and women for demographics, BMI, obese status, CESD score, or depression classification. Men and women reported equal engagement (29%) in at least one unhealthy weight loss practice. Only 14% of the sample engaged in more than two unhealthy weight loss practices and less than 5% in more than three. The most common unhealthy weight loss practices were "eating very little" and "skipping meals." College students engaging in at least one unhealthy weight loss practice had a statistically greater CESD score ( $M = 7.88$ ,  $SD = 3.98$ ) than college students not engaging in any ( $M = 5.29$ ,  $SD = 4.84$ ),  $t(402) = 5.59$ ,  $p = < 0.001$ .

Download English Version:

<https://daneshyari.com/en/article/906493>

Download Persian Version:

<https://daneshyari.com/article/906493>

[Daneshyari.com](https://daneshyari.com)