

Original research article

Blood copper levels in Mexican users of the T380A IUD

Dolores De la Cruz^a, Angeles Cruz^a, Maricela Arteaga^b, Lourdes Castillo^b, Horacio Tovalín^{c,*}

^aUnidad Universitaria de Investigación en Cariología (University Cariology Research Unit), FES-Zaragoza, Universidad Nacional Autónoma de México, Mexico City, Mexico

^bLaboratorio de Espectroscopía (Spectroscopy Laboratory), FES-Zaragoza, Universidad Nacional Autónoma de México, Mexico City, Mexico

^cDivisión de Estudios de Posgrado e Investigación (Graduate Studies and Research Division), FES-Zaragoza, Universidad Nacional Autónoma de México, Mexico City, Mexico

Received 23 November 2004; revised 15 February 2005; accepted 17 February 2005

Abstract

Objective: Copper T380A is one of the most commonly prescribed intrauterine devices (IUDs). However, there are few reports of its systemic health effects. This study evaluates the association between the use of T380A IUD and blood Cu levels in a group of Mexican users.

Methods: Eighty-six T380A users and eight nonusers were recruited. Blood Cu levels were measured using an atomic absorption spectrophotometric technique.

Results: In IUD users, mean blood Cu level was significantly higher than in nonusers, 216.63 vs. 107.47 $\mu\text{g/dL}$ ($p \leq .001$), and above the normal blood Cu level range of 80–160 $\mu\text{g/dL}$. No significant difference in blood Cu levels was observed by age or length of time of IUD use.

Conclusions: The observed higher blood Cu levels among T380A user stands out in sharp contrast to previous reports of other types of Cu IUDs. Further research should evaluate if the observed levels are associated to toxic effects with the general population or special groups.

© 2005 Elsevier Inc. All rights reserved.

Keywords: Copper; T380A; Blood; Users; Mexican

1. Introduction

Copper intrauterine devices (Cu IUDs) are valuable birth control resources. Release of Cu from these devices is necessary for their contraceptive effects [1]. The Cu T380A is one of the most commonly prescribed IUDs; some of its advantages are fewer side effects in comparison with other method [2]. However, there are few reports of the T380A systemic side effects. The T380A is made of plastic with added barium sulfate for X-ray visibility, with a T-shape 36-mm height and 32-mm width. A copper wire twisted on its body and two copper sleeves on its arms, with a total Cu contact surface of 380 mm² [2].

Release and absorption rates have been evaluated for different Cu IUDs. A study carried out by Hagenfeldt [3] did not find a significant difference in plasma Cu concentrations levels between Cu T120 and T130 users and nonusers, but observed an increment in endometrial Cu concentration. The same author [3] estimated a T120 and T135 annual Cu liberation of 10.3 mg/year (range=7.4–13.5 mg/year) and a

5% annual endometrial Cu absorption (about 0.5 mg per year), and for T200, a daily release of 45 μg , around 18 mg/year. Timonen [4] reported a daily release that ranged from 26 to 74 μg Cu/day for different IUDs, and Thiery and Kosonen [5] reported a release of 23 μg Cu/d after 10 years of insertion for T220 devices without calcareous deposits. In 1972, Oster [6] observed that Cu ions in uterus could cross over to blood, increasing about 1% blood Cu concentration in T200A users. Later, Anteby et al. [7] found a nonstatistical difference between the serum levels of Cu before and after insertion of the T200A IUD. Prema et al. [8] studied 24-h urinary Cu excretion and did not find differences between a control group and Cu IUD users.

Copper is an essential trace element found in small amounts in different cells and tissues. Copper exists in both oxidized, cupric (Cu^{2+}) or reduced (Cu^+) state [9]. Copper's function as a cofactor is required for structural and catalytic properties of a variety of important enzymes, such as cytochrome *c* oxidase, Cu-zinc superoxide dismutase, tyrosinase, dopamine beta-hydroxylase and others. Once Cu is absorbed, it is stored in several tissues and organs, among which, liver, kidney, brain and bone have the largest amount and blood the lowest [10]. Bile is the main route for Cu

* Corresponding author. Tel.: +52 55 56 23 07 21.

E-mail address: htovalin@hotmail.com (H. Tovalín).

Table 1
Blood Cu levels in T380A users and nonusers

Group	Blood Cu, µg/dL (µmol/L)				
	N	Mean	SD	Minimum	Maximum
Users	86	217.36 ^a (34.12)	±21.32	178 (27.95)	277 (43.49)
Nonusers	8	107.47 (16.87)	±16.30	89 (13.97)	134 (21.03)

^a *t* Test, *p*≤.001.

excretion, while urine represents a minor route [11,12]. The daily recommended Cu intake is 1.5–3 mg/day [13,14]. Food Cu concentrations are variable, chocolate and liver being among the highest in concentration, followed by cereals and fruits.

In the general population, a mean serum Cu concentration of 80 µg/dL with an upper range of 150 µg/dL [15,16] and a whole blood Cu concentration with a mean value of 110 µg/dL have been reported [16,17]. Although Cu is an essential element, it can become toxic for cells when present at high concentrations; as it is the case with cadmium and lead, Cu could displace metal ions such as calcium and magnesium [18]. In addition, the Cu²⁺ as other redox active metals can be toxic by generating free hydroxyl radicals [19,20]. Free hydroxyl radicals resulting from catalytic Cu can bind to DNA and result in DNA strand breaks and DNA base modifications [21].

Toxic symptoms such as gastrointestinal symptoms have been reported at a serum level of 200 µg/dL and whole blood concentration above 287 µg/dL [16]. Hepatic symptoms appear at blood levels above 798 µg/dL [22]. Very high Cu levels can cause acute toxicity, and an oral dose of 200 mg/kg/day is considered fatal [23]. Individuals with Wilson's disease, an autosomal recessive disease, do not adequately metabolize and eliminate Cu and have a progressive accumulation of Cu in the brain, kidneys and liver [24].

In an effort to provide more information on Cu absorption in T380A IUD users, a group of users and nonusers was studied with the aim to measure their blood Cu levels and to evaluate potential Cu toxic effects.

2. Material and methods

The study was approved by the Research Committee of the Facultad de Estudios Superiores-Zaragoza, Universidad Nacional Autónoma de México (UNAM).

2.1. Study population

Volunteers were recruited among women who attended a general hospital located in Mexico City. Cases were healthy women 15–45 years old who have been using the T380A IUD for at least 1 month. Controls were healthy women from the same age group not using any type of birth control method. Verbal and printed information about the study was provided to the hospital clients, and those who accepted to participate received and signed an informed consent. Women taking any kind of medication or vitamin supplements were excluded from this study.

2.2. Blood sample collection and analysis

Blood was collected in heparinized sample vials by a nurse and kept in refrigeration until analysis. Blood samples were processed as described by ACF [25]. Briefly, blood was lysed with 1% Triton X-100 solution (Sigma-Aldrich, St. Louis, MO, USA). Blood Cu was chelated with ammonium pyrrolidine dithiocarbamate (Sigma-Aldrich). Copper chelates were extracted with methyl isobutyl ketone (Merk & Co., Whitehouse Station, NJ, USA), and analyzed by an atomic absorption spectrometer Perkin-Elmer 3110 (Perkin-Elmer, Boston, MA, USA).

2.3. Data collection and statistical analysis

A questionnaire about demographics, length of time of IUD use, diet and health information was administered by trained interviewers. Length of time of IUD use was measured by counting the number of months since IUD insertion. Age and IUD use subgroups were created using the values of the first, second and third quartiles as cut-points.

Student's *t* test was used to compare mean values of cases and controls, while ANOVA was used to compare blood Cu levels among age subgroups and to compare the Cu levels among subgroups of length of IUD use. The statistical software SPSS 10 (SPSS, Chicago, IL, USA) was used to analyze the data.

3. Results

Ninety-four women participated in the study, 86 were T380A users and 8 were controls. The mean ages for the cases and controls were 27.41±6.62 and 23.2±4.34 years old, respectively. No difference between the groups was found related to consumption of foods with relatively high Cu content such as chocolate, cereals, vegetables and fruits. Mean length of IUD use was 25±1.12 months with a minimum of 2 months and a maximum of 124 months.




Fig. 1. Blood Cu levels (µg/dL) in T380A users by age group. *CU mean ± SD µg/dL (µmol/L) by age groups: 15–22=217.06 (34.07)±21.85; 23–25=226.73 (35.60)±26.30; 26–30=215.09 (33.77)±14.87; 31–39=211.25 (33.17)±19.36.

Download English Version:

<https://daneshyari.com/en/article/9317201>

Download Persian Version:

<https://daneshyari.com/article/9317201>

[Daneshyari.com](https://daneshyari.com)