

A role of insulin-like growth factor 1 in β amyloid-induced disinhibition of hippocampal neurons

Changhong Xing^{a,**}, Yanling Yin^{b,**}, Rui Chang^a, Xiangping He^a, Zuoping Xie^{a,*}

^a Department of Biological Science and Biotechnology, State Key Laboratory of Biomembrane and Membrane Biotechnology, Tsinghua University, Beijing 100084, China

^b Department of Neurobiology, The Capital University of Medical Sciences, Beijing 100054, China

Received 23 February 2005; received in revised form 1 April 2005; accepted 17 April 2005

Abstract

In the present study we investigated the effects of β amyloid ($A\beta$) on inhibitory synaptic transmission in the cultured hippocampal neurons using whole-cell patch-clamp recordings and immunocytochemistry, and examined the role of insulin-like growth factor 1 (IGF-1). Incubation with 4 μ M $A\beta_{25-35}$ for 24 h significantly decreased the frequency of spontaneous inhibitory postsynaptic currents (sIPSCs), but had no effect on the mean amplitude. Pretreatment with 10 ng/ml IGF-1 for 24 h prior to $A\beta_{25-35}$ exposure blocked $A\beta$ -induced disinhibition of hippocampal neurons. The frequency and mean amplitude of miniature IPSC (mIPSCs) were not significantly affected by $A\beta$. The rise and decay kinetics of sIPSCs and mIPSCs were similar for the control and $A\beta_{25-35}$ -treated hippocampal neurons. Immunocytochemistry showed no changes in the ratio of γ -aminobutyric acid (GABA) positive cells subsequent to treatment with $A\beta$, or IGF-1. Together these data suggest that $A\beta$ -induced the disinhibition in cultured hippocampal neurons, whereas IGF-1 could block this effect.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Insulin-like growth factor 1; β Amyloid; Inhibitory postsynaptic current; Whole-cell patch-clamp; γ -Aminobutyric acid

Alzheimer's disease (AD) is a progressive neurodegenerative disease and the most frequent cause of cognitive deficit in the aged. Insoluble fibril deposits of $A\beta$ is the major component of senile plaques and plays an important role in the neurodegenerative process [11,16,20]. It is well established that the cholinergic and glutamatergic systems are adversely affected in the progressive course of AD. However, other neurotransmitter systems, such as serotonergic [1,7,15], histaminergic [18], and GABAergic [17] systems are less well studied. Previous studies have focused on the effects of $A\beta$ on excitatory synaptic transmission but little is known how $A\beta$ influences inhibitory synaptic transmission. Therefore, to elucidate the effects of $A\beta$ on this important pathway, we recorded sIPSCs and miniature IPSCs using whole-cell patch-clamp in cultured rat hippocampal neurons, and counted the numbers of GABA positive cells using immunocytochemical staining.

IGF-1 is a pleiotropic factor with structural and functional homologies to IGF-2 and insulin. IGF-1 has important functions in the brain, including metabolic, neurotrophic, neuro-modulatory and neuroendocrine actions [19]. IGF-1 and its receptor are highly concentrated in the hippocampus, an area severely affected in AD [6,9]. Recently, IGF-1 has gained increasing attention for the pathogenesis of age-related neurodegenerative diseases [12]. AD patients show changes in insulin and IGF-1 levels, which may protect hippocampal neurons against the toxicity of $A\beta$ [8]. Additionally, IGF-1 may be a key factor in regulating the clearance of $A\beta$ from the brain [4]. Based on these previous reports, we investigated the effects of IGF-1 in the regulation $A\beta$ -induced changes of inhibitory synaptic transmission.

Hippocampal neuron cultures were prepared as described previously [10]. Briefly, pregnant Wistar rats were anesthetized with pentobarbital and the E18–19 embryos delivered by cesarean section. The hippocampi were dissected and incubated with 0.25% Trypsin–EDTA for 15 min at 37 °C and mechanically dissociated. The resulting single cell

* Corresponding author. Tel.: +86 10 62788677.

** Co-corresponding author.

E-mail address: zuopingx@mail.tsinghua.edu.cn (Z. Xie).

suspension was diluted at a density of 1×10^5 cells/ml in high glucose DMEM containing 10% fetal bovine serum, 5% equine serum and 2 mM L-glutamine, then plated in 35 mm-cell plates coated with poly-D-lysine (20 μ g/ml). Cells were incubated at 37 °C in a humidified incubator with 5% CO₂. After approximately 20 h, the medium was replaced by serum-free Neurobasal medium containing B27 supplement and 0.5 mM L-glutamine to inhibit the growth of glia cells. Every 3–4 days half of the media was replaced and the cultures were used for experiments on 10–14 days after plating. A β 25–35 (Sigma) was dissolved in sterile distilled water at a concentration of 2 mM as a stock solution and incubated at 37 °C for 96 h before use. To determine the effects of A β , we incubated 4 μ M A β 25–35 with cultured hippocampal neurons for 24 h and measured electrophysiological and immunocytochemical changes. In order to explore the action of IGF-1 on these effects, cultures were pretreated with 10 ng/ml IGF-1 (PeproTech) for 24 h prior to exposure to A β 25–35. Data were expressed as the mean \pm S.E.M. Statistical significance was determined as $p < 0.05$ by one-way ANOVA (SPSS, Chicago, IL). In all cases, n refers to the number of neurons studied from multiple dissections and data pooled.

The whole-cell patch-clamp technique was used to record current. The patch electrodes of thick-walled boro-silicate glass (VWR Scientific) were pulled on a PP-83 micropipette puller (Narishige). The patch-pipette solution contained (in mM): 140 KCl, 10 HEPES, 10 EGTA, 2 MgCl₂, 2 Na₂ATP, 1 CaCl₂, pH 7.3. The typical resistance of glass electrodes was 3–5 M Ω when filled with intracellular pipette solution. The range of the whole-cell series resistance is 10–15 M Ω . Data were collected with an Axopatch 200B amplifier (Axon Instruments) and acquired and analyzed using pCLAMP 9 (Axon Instruments). During experiments, culture dishes were rinsed and perfused with extracellular solution containing (in mM) 140 NaCl, 5 KCl, 1 MgCl₂, 0.5 CaCl₂, 10 Glucose, 10 HEPES, pH 7.4. Synaptically spontaneous IPSCs were isolated by the application of 20 μ M 6,7-dinitroquinoxaline-2,3-dione (DNQX) and 50 μ M aminophosphonobutyrate (APV), antagonist to the excitatory α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) and *N*-methyl-D-aspartate (NMDA) receptors, respectively. mIPSCs were recorded using 1 μ M TTX added to the extracellular solution. sIPSCs and mIPSCs were recorded without synaptic stimulation at a holding potential of -70 mV from cultured hippocampal neurons for at least 5 min. The currents of sIPSCs were completely abolished after using 50 μ M bicuculline, the antagonist of the inhibitory GABA_A receptor, indicating that they were mediated by GABA_A receptors. Cells were chosen for recording based on their morphology and density of surrounding cells. Relatively isolated or spherically shaped cells were avoided.

A β 25–35 treatment for 24 h significantly decreased the frequency of sIPSCs in cultured hippocampal neurons, but had little influence on the mean amplitude of sIPSCs (Fig. 1). Quantitative analysis of all neurons recorded indicated that sIPSCs frequency decreased by 47.5%. The average value of

frequency of the normal group ($n = 10$) was 3.37 ± 0.33 Hz. However, after treatment with 4 μ M A β 25–35 for 24 h, the frequency decreased to 1.77 ± 0.43 Hz ($n = 11$). The similar results were observed with A β 1–42. After incubation with 250 nM A β 1–42 for 24 h, the frequency of sIPSCs decreased to 2.08 ± 0.45 Hz ($n = 12$), and the mean amplitude did not change. A β 25–35 treatment had no significant effects on both the frequency and mean amplitude of mIPSCs (Fig. 2). When the neurons were pretreated with 10 ng/ml IGF-1 for 24 h prior to exposure to A β 25–35, some inhibitory firings were saved and the frequency of sIPSCs was 2.95 ± 0.36 Hz ($n = 10$). There were no significant difference between the control and IGF-1 pretreated group (Fig. 1). Treatment with IGF-1 alone did not affect the frequency of sIPSCs. Therefore, it appears that A β 25–35-induced synaptic disinhibition and this effect could be blocked by IGF-1.

Furthermore, the other postsynaptic properties of sIPSCs and mIPSCs including rise time constant, decay time constant and the time to peak were analyzed using Clampfit 9.0 (Axon Instruments). The rise and decay kinetics of sIPSCs and mIPSCs were similar for the control and A β 25–35-treated hippocampal neurons, and there are no significant differences between the two groups. The details are in Table 1.

The numbers of GABAergic neurons was examined in cultured hippocampal neurons using immunocytochemistry. Hippocampal neurons were fixed with 4% paraformaldehyde for 30 min at room temperature, permeabilized with 0.1% Triton X-100 for 5 min, and blocked with 5% horse serum for 30 min at 37 °C. The cultures were incubated with rabbit anti-GABA primary antibody (Sigma, 1:1000) for 1 h at 37 °C and followed by TRITC-labeled secondary antibody (1:200) for 45 min at room temperature. Cells were examined using a fluorescence microscope with six to ten high magnification (400 \times) fields randomly selected for counting. Experiment was repeated three times.

A β 25–35 treatment did not affect the ratio of GABA-positive cells (Fig. 3). In the control group, the percentage of GABA positive cells was 30.19%. After incubated with 4 μ M of A β 25–35 for 24 h, GABA positive cells were 28.99%. Pretreatment with IGF-1 prior to exposure to A β 25–35 or treatment with IGF-1 alone did not change the ratio.

GABA mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel. In addition, GABA functions as an excitatory modulator: it depolarizes the membrane of embryonic neurons and axons, exerts trophic, chemoattractant or chemokinetic influences that promote cell migration to form specific systems [2,5]. A β -induced lesions are involved in decreasing GABAergic transmission. It has been reported that GABA and the GABA_A receptor agonists protect neurons against A β -induced neurotoxicity [13]. In the current study, A β significantly reduced the frequency of sIPSCs but did not affect the frequency of mIPSCs, indicating that the effects of A β on inhibitory synaptic transmission are action potential-related (i.e., TTX-sensitive). To further clarify the action site of A β -induced disinhibition in hippocampal

Download English Version:

<https://daneshyari.com/en/article/9429241>

Download Persian Version:

<https://daneshyari.com/article/9429241>

[Daneshyari.com](https://daneshyari.com)