



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Scientia Horticulturae 106 (2005) 125–128

SCIENTIA
HORTICULTURAE

www.elsevier.com/locate/scihorti

Short communication

Susceptibility of Gisela 5 and Maxma 14 cherry rootstocks to four *Phytophthora* species

E. Exadaktylou, T. Thomidis*

National Agriculture Research Foundation (NAGREF), Pomology Institute,
S.S. Naoussas 38, P.C. 59200 Naoussa, Greece

Received 29 November 2004; received in revised form 10 February 2005; accepted 10 February 2005

Abstract

Susceptibility of Gisela 5 and Maxma 14 cherry rootstocks to *Phytophthora cactorum*, *P. citrophthora*, *P. citricola* and *P. parasitica* was evaluated. All *Phytophthora* species were pathogenic to both of the cherry rootstocks tested. The level of susceptibility, however, varied according to *Phytophthora* species. *P. citrophthora* and *P. parasitica* isolates were the most virulent. *P. citricola* showed moderately virulence. The least virulence was shown from the *P. cactorum* isolates. Both rootstocks showed similar susceptibility.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Cherry; Crown rot; *Phytophthora*; Rootstocks

1. Introduction

Gisela 5 and Maxma 14 cherry rootstocks are among the best dwarfing, precocious and productive rootstocks for modern, intensive sweet cherry growing (Zimmermann, 1994). In the last few years, Gisela 5 and Maxma 14 have replaced *Prunus avium* and Mazzard rootstocks in Greece because of their ability to produce dwarfing and precocity. These rootstocks show very good adaptation to Greek soil-climate conditions. Crown rot, caused by *Phytophthora* spp., is a serious disease of cherry trees (Erwin and Ribeiro, 1996). In Greece, *P. citrophthora*, *P. cactorum*, *P. citricola*, *P. parasitica* and *P. syringae* were

* Corresponding author. Tel.: +30 233 204 1548; fax: +30 233 204 1178.
E-mail address: thomi-1@otenet.gr (T. Thomidis).

reported as causal agents of crown rot diseases on cherry trees (Thomidis, 2001; Thomidis and Sotiropoulos, 2003).

The aim of this study was to evaluate the susceptibility of Gisela 5 and Maxma 14 cherry rootstocks to four *Phytophthora* species (*P. cactorum*, *P. citrophthora*, *P. citricola*, *P. parasitica*).

2. Materials and methods

Six isolates of *P. cactorum* (BPIC-1168, BPIC-1128, both isolated from the trunk of infected almond trees) *P. citrophthora* (BPIC-1262 isolated from the trunk of citrus, BPIC-1267 isolated from the trunk of a pear tree), *P. parasitica* (BPIC-1263 isolated from the trunk of citrus) and *P. citricola* (BPIC-1908 isolated from lemon fruit) were used in this study.

In an excised twig laboratory assay (Jeffers et al., 1981), cornmeal agar amended with antibiotics (primaricin 10 mg, ampicillin 250 mg, rifampicin 10 mg) was dispensed in sterilized jars to a depth of 10 mm. Two agar plugs with mycelium of a tested isolate were placed into each jar. Jars were sealed and placed in incubators at 25 °C in the dark. Woody shoots, about 1 cm diameter and 7 cm long, were collected from 2-year-old mother trees. Using a flamed sharp knife, segments were trimmed to a slant at the base. Two jars for each *Phytophthora* isolate were used to inoculate 20 segments of each cherry rootstock. Jars were then resealed and returned to incubators for 4 days. By subtracting of the depth of agar from the total length of necrosis a value for necrosis was obtained. A second laboratory experiment was conducted using an excised shoot assay (Matheron and Mircetich, 1985). Segments of woody shoot of each rootstock, 6 cm in length and 1.5–2 cm in diameter were collected from the same mother trees. Thirty segments (10 from each of the three tested rootstocks) were inoculated with each isolate of *Phytophthora*. The inoculum, which consisted of a 6 mm in diameter plug from a 5-day-old culture, was inserted in the middle of excised shoot pieces under the bark. The wound was covered with petroleum jelly and wrapped with adhesive tape to prevent desiccation. Inoculated shoot segments were incubated for 4 days at 25 °C in moist chambers, after which the length of the resulting necrosis was recorded. A field experiment was conducted using a rootstock assay (Thomidis, 2001). Two-year-old trees were used in these experiments. Inoculations were performed on the trunk, about 10 cm above soil surface, by transferring an agar plug containing mycelium of the pathogen onto a wound. Each wound was then covered with petroleum jelly and wrapped with adhesive tape to prevent desiccation. Fifteen days later, the length of resulting necrosis was measured. Twenty plants (10 of each rootstock) were inoculated with each *Phytophthora* isolate. Temperatures during experiments ranged from 18 to 28 °C. P₅ARP selective medium was used for re-isolation of fungi (Jeffers and Martin, 1986).

3. Results and discussion

Results from both laboratory and field experiments are presented in Table 1. All isolates were pathogenic on Gisela 5 and Maxma 14 cherry rootstocks. In laboratory experiments, the isolates BPIC-1263 (*P. parasitica*), BPIC-1262 (*P. citrophthora*) and BPIC-1267 (*P.*

Download English Version:

<https://daneshyari.com/en/article/9488823>

Download Persian Version:

<https://daneshyari.com/article/9488823>

[Daneshyari.com](https://daneshyari.com)