

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SciVerse ScienceDirect

journal homepage: [www.elsevier.com/locate/jval](http://www.elsevier.com/locate/jval)

## Using a Survey to Estimate Health Expectancy and Quality-Adjusted Life Expectancy to Assess Inequalities in Health and Quality of Life

Brendan Collins, M.Sc.\*

University of Liverpool/NHS Wirral, Merseyside, UK

### ABSTRACT

**Background:** There has been a policy debate in the United Kingdom about moving beyond traditional measures of life expectancy and economic output to developing more meaningful ways of measuring national well-being. **Objective:** To test whether quality adjusted life expectancy (QALE) was a useful indicator of health inequalities. **Methods:** EuroQol five-dimensional questionnaire data from a well-being survey was combined with actuarial life expectancy (LE) data to estimate healthy LE (HLE), that is, years of life lived in good health, and QALE, that is, quality-adjusted life-years (QALYs) lived for Wirral, a borough in the north west of England. **Results:** The gap between Wirral and the most deprived areas was 4.45 years for LE, 5.34 for QALE, and 7.55 for HLE. The gap in QALE was 20% greater than the gap in LE, while the gap in HLE was 70% greater. **Conclusions:** The fact that the QALE gap value lies between the HLE value and the LE value

suggests that QALE is a more sensitive indicator than HLE, as in this study QALE is derived from 243 possible EuroQol five-dimensional questionnaire profiles whereas HLE is based only on whether or not an individual rates his or her health as good, a binary variable. This study discusses how QALE could be a useful indicator for measuring health inequalities in future, especially as cost utility and QALYs are seen as the gold standard used by the National Institute for Health and Clinical Excellence in the United Kingdom to measure outcomes for health interventions in England, and discusses how a monetary valuation of QALYs could be used to put a societal cost on health inequalities. **Keywords:** EQ-5D, health inequalities, population surveys, QALYs.

Copyright © 2013, International Society for Pharmacoeconomics and Outcomes Research (ISPOR). Published by Elsevier Inc.

### Introduction

In the United Kingdom, there has been a recent policy debate about regarding well-being as an economic good, measured alongside established measures of income, such as gross domestic product, and health, such as life expectancy (LE) [1]. This change in focus chimes with Organisation for Economic Co-operation and Development's Istanbul Declaration [2] on improving well-being and considers the Easterlin paradox first described in 1974 [3]—that increasing income does not always increase happiness, and hedonic treadmill theory, that adverse life events do not change an individual's level of happiness as much as expected [4]. The UK Office for National Statistics (ONS) has formulated well-being measures across 10 domains: the economy, individual well-being, our relationships; where we live, health, natural environment, personal finance, what we do, governance, and education and skills [5].

The United Kingdom had increasing levels of income inequalities since the 1970s, with inequalities in health outcomes remaining despite targeted investment [6]. The gap in health expectancy or healthy LE (HLE) between areas is typically wider than the gap in LE, indicating that health inequalities are greater when morbidity and mortality are combined. In the EU-27 countries, the largest LE gap between countries for males is 12.3 years (between Iceland and Lithuania) whereas the largest HLE gap is 50% greater at 18.4

years (between Sweden and Slovakia). For females, the largest LE gap is 7.6 years whereas the largest HLE gap is 18.3 years (data for 2009 [7]). A study comparing quality-adjusted LE (QALE) across countries found some interesting patterns, with women in two countries (Spain and The Netherlands) having a smaller QALE gap than the LE gap, meaning that Spanish women live longer with more health problems than Dutch women [8].

The ONS has previously measured disability-free LE as well as HLE at birth and at age 65 years, calculated by combining actuarial cohort LE data with survey data. Although the EuroQol five-dimensional (EQ-5D) questionnaire is used in population health surveys such as the Health Survey for England, it has not been routinely used to assess quality-adjusted life-years (QALYs) experienced across a population. There is a disparity between the UK gold standard in measuring health outcomes (the EQ-5D questionnaire and QALYs recommended by the National Institute for Health and Clinical Excellence) and what is seen as the gold standard in measuring health status across the population (measures such as the HLE recommended by the ONS). Internationally, disability-adjusted life-years are used for the World Health Organization's Global Burden of Disease project, which was recently updated [9].

The EQ-5D 3-level questionnaire is a self-reported health-related quality of life tool that consists of five dimensions (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression) each of which can take one of three levels of

\* Address correspondence to: Brendan Collins, University of Liverpool and NHS Wirral, Old Market House, Hamilton Street, Birkenhead, Merseyside CH41 5AL, UK.

E-mail: [Brendan.collins@nhs.net](mailto:Brendan.collins@nhs.net).

1098-3015/\$36.00 – see front matter Copyright © 2013, International Society for Pharmacoeconomics and Outcomes Research (ISPOR).

Published by Elsevier Inc.

<http://dx.doi.org/10.1016/j.jval.2013.01.004>

severity (no problems/some or moderate problems/extreme problems). The EQ-5D questionnaire profiles are matched to UK utility scores, giving the desirability of a particular health state, measured between  $-0.594$  (worst health state) and  $1$  (perfect health). In a randomized controlled trial, change in utility as a result of a health intervention is measured in the same person at baseline and at set time intervals, so that any change can be attributed to the intervention. This change in utility is used to calculate QALYs experienced. The EQ-5D questionnaire being self-reported has an element of subjectivity where individuals may have similar health status but responses indicate different levels of health problems. In a randomized controlled trial, individual improvement in the EQ-5D questionnaire is used to calculate the QALYs gained, and so this accounts for some of the subjectivity, an improvement is always an improvement. But in a population-level study such as this, each individual is completing the EQ-5D questionnaire once; however, with a large sample size (1522 people in this study), some of these subjective differences would even out across the population. This element of subjectivity is also true for HLE, which is widely used as a measure of health status. It has been claimed that the EQ-5D questionnaire is not sensitive in measuring health problems such as fatigue, sensory impairment, or mental health problems, and if so then the impact of these conditions would be underrepresented in QALE derived from EQ-5D questionnaire survey data.

The aim of this study was to show that because QALE is based on the EQ-5D questionnaire profile, which has 243 possible health states, QALE will be more robust as an indicator of population health than LE or HLE, which are both essentially based on binary variables, that is, whether after a period of time an individual is still alive, and if he or she is, whether he or she rates his or her health as good.

## Methods

LE, HLE, and QALE were calculated for Wirral, a borough in the northwest of England, with an estimated ONS population of 309,000 people in 2009. This area was chosen because Wirral has extremes of affluence and poverty, with the east side containing some of the most deprived areas in England and the west side being an affluent retirement destination. Data were combined from a well-being survey [10] that was commissioned for the northwest of England ( $N = 1522$  for Wirral), and carried out in

2009, and mortality and population data for 2005–2007 (3 years pooled), the most recent data available when the results were analyzed. The methods for collecting the survey are described in more detail elsewhere [11]. Survey data were weighted by age, gender, and deprivation, and so the scores should represent a true average. The weighted EQ-5D questionnaire index scores and health status scores were combined for males and females and grouped into six age bands, 16 to 17, 18 to 24, 25 to 39, 40 to 54, 55 to 64, and 65 years and older. These utility scores used the UK EQ-5D questionnaire value set produced by EuroQol using a representative sample (3359 people) of the UK population using the time trade-off method [12]. Because the well-being survey was carried out only on individuals aged 16 years and older, a maximum utility score of 1 and a probability of reporting oneself as being healthy of 1 was assumed for those younger than 16 years.

Cohort LE was calculated by using the Chiang II method [13] used by the UK ONS. The utility and self-reported health data were combined with the LE data by using the method outlined by Sullivan in 1971 [14]. This is where QALE is calculated as follows:

$$QALE = \frac{\sum_a (U_a \cdot P_a)}{\sum P} \cdot LE$$

where  $U_a$  is average utility in age group  $a$ ,  $P_a$  is the population surviving in age group  $a$ ,  $z$  is the maximum age group, and LE is total cohort life expectancy (years).

To understand inequalities in health and quality of life, the analysis was carried out for the whole of Wirral, as well as for the areas of Wirral that fell into the 20% most deprived and 20% least deprived lower layer super output areas (a small area geography used by the ONS, where each contains on average 1500 people) nationally based on the Index of Multiple Deprivation 2007, which is a widely used UK deprivation measure [15]. Of the Wirral population at the time, 32% fell into the most deprived quintile and 10% into the least deprived quintile.

## Results

The differences in utility and LE were analyzed for Wirral as a whole and for the most and least deprived quintiles. The characteristics of respondents from each group are shown in Table 1. The least deprived areas have a greater proportion of males answering the survey, are older on average, and have a greater proportion of people in employment. The groups were similar for average mental well-being score as measured by using

**Table 1 – Comparative statistics for the most and least deprived areas and the whole of Wirral.**

	Most deprived	Least deprived	Whole of Wirral
N	687	75	1522
Average age (y)	49.6	57.9	52.4
Gender, male (%)	36.3	48	38.9
Average Short Warwick-Edinburgh Mental Wellbeing Scale score	27.75	28.09	28.00
Work status (%)			
Not recorded	0.9	1.3	1.2
Full-time education	2.3	1.3	1.6
Not working for domestic reasons	9.8	0.0	6.8
Other	1.7	0.0	1.3
Out of work, registered unemployed but not actively seeking work	2.6	1.3	2.2
Out of work, registered unemployed and actively seeking work	8.3	4.0	6.8
Paid work: full-time	23.6	41.3	25.8
Paid work: part-time	7.6	10.7	9.2
Permanently sick or disabled	10.9	0.0	7.5
Refused	0.4	0.0	0.5
Retired	30.4	36.0	35.2
Self-employed	1.5	4.0	2.0

Download English Version:

<https://daneshyari.com/en/article/989323>

Download Persian Version:

<https://daneshyari.com/article/989323>

[Daneshyari.com](https://daneshyari.com)