

The activity of brain and liver cytochrome P450 2D (CYP2D) is differently affected by antidepressants in the chronic mild stress (CMS) model of depression in the rat

Anna Haduch, Marta Rysz, Mariusz Papp, Władysława A. Daniel

PII: S0006-2952(18)30387-3
DOI: https://doi.org/10.1016/j.bcp.2018.09.005
Reference: BCP 13284

To appear in: *Biochemical Pharmacology*

Received Date: 5 July 2018
Accepted Date: 5 September 2018

Please cite this article as: A. Haduch, M. Rysz, M. Papp, W.A. Daniel, The activity of brain and liver cytochrome P450 2D (CYP2D) is differently affected by antidepressants in the chronic mild stress (CMS) model of depression in the rat, *Biochemical Pharmacology* (2018), doi: https://doi.org/10.1016/j.bcp.2018.09.005

The activity of brain and liver cytochrome P450 2D (CYP2D) is differently affected by antidepressants in the chronic mild stress (CMS) model of depression in the rat

Anna Haduch[1], Marta Rysz[1], Mariusz Papp[2], Władysława A. Daniel[1]*

[1]*Department of Pharmacokinetics and Drug Metabolism, Institute of Pharmacology, Polish Academy of Sciences, Smętna 12, 31-343 Kraków, Poland*

[2]*Department of Pharmacology, Institute of Pharmacology, Polish Academy of Sciences, Smętna 12, 31-343 Kraków, Poland*

*Corresponding author. Tel.: +48-12-6623266; fax: +48-12-6374500; e-mail address: nfdaniel@cyf-kr.edu.pl

Abbreviations: ACTB, beta-actin; CYP, cytochrome P450; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; CMS, chronic mild stress; HPA, hypothalamic-pituitary-adrenal axis; MD, early-life maternal deprivation; RS, repeated restraint stress; SSRI, serotonin selective reuptake inhibitor; ESC, escitalopram, VEN, venlafaxine