

## Accepted Manuscript

Impact of two light programs and two levels of dietary tryptophan for weanling piglets

Lívea Maria Gomes , Gabriela de Mello Miassi ,  
Luan Sousa dos Santos , Mayra Anton Dib Saleh ,  
José Roberto Sartori , Marcos Lívio Panhoza Tse ,  
Dirlei Antonio Berto

PII: S1871-1413(18)30259-2  
DOI: <https://doi.org/10.1016/j.livsci.2018.08.010>  
Reference: LIVSCI 3517



To appear in: *Livestock Science*

Received date: 22 September 2017  
Revised date: 15 August 2018  
Accepted date: 22 August 2018

Please cite this article as: Lívea Maria Gomes , Gabriela de Mello Miassi , Luan Sousa dos Santos , Mayra Anton Dib Saleh , José Roberto Sartori , Marcos Lívio Panhoza Tse , Dirlei Antonio Berto , Impact of two light programs and two levels of dietary tryptophan for weanling piglets, *Livestock Science* (2018), doi: <https://doi.org/10.1016/j.livsci.2018.08.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- The photoperiod and Trp in diet affected plasma cortisol and piglet behavior
- Piglets that received the highest Trp level had lower plasma cortisol levels
- Feed:gain ratio improved for piglets receiving 12-h light/day
- 23-h of light/day indicated a higher stress level of animals

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/9954656>

Download Persian Version:

<https://daneshyari.com/article/9954656>

[Daneshyari.com](https://daneshyari.com)