

Accepted Manuscript

Treatment of intraurethral condylomata acuminata with surgery and cidofovir instillations in two immunocompromised patients and review of the literature.

H.J. Florin, R. Snoeck, B. Van Cleynenbreugel, M. Albersen



PII: S0166-3542(18)30396-6

DOI: [10.1016/j.antiviral.2018.08.016](https://doi.org/10.1016/j.antiviral.2018.08.016)

Reference: AVR 4362

To appear in: *Antiviral Research*

Received Date: 27 June 2018

Revised Date: 14 August 2018

Accepted Date: 24 August 2018

Please cite this article as: Florin, H.J., Snoeck, R., Van Cleynenbreugel, B., Albersen, M., Treatment of intraurethral condylomata acuminata with surgery and cidofovir instillations in two immunocompromised patients and review of the literature., *Antiviral Research* (2018), doi: 10.1016/j.antiviral.2018.08.016.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Treatment of intraurethral condylomata acuminata with surgery and cidofovir instillations in two immunocompromised patients and review of the literature.

HJ. Florin^a, R. Snoeck^b, B. Van Cleynenbreugel^a, M. Albersen^a

^a Department of Urology, University Hospitals Leuven, Leuven, Belgium.

^b Rega Institute for Medical Research, KU Leuven, Belgium.

Corresponding author

Florin Hendrik-Jan

hjflorin@gmail.com

department of urology UZ Leuven

Herestraat 49

3000 Leuven

Belgium

Abstract

Condylomata acuminata (CA) or anogenital warts are benign proliferative lesions caused by low-risk human papillomaviruses (HPV). Treating CA can be very frustrating for patients and clinicians due to the high recurrence rates. Immunosuppression is associated with larger size of CA that are more frequently resistant to treatment. Surgical approaches tend to be poorly effective in the long-term because of high recurrence rates related to the persistence of HPV-infected cells. In our search to find an agent to treat intraurethral CA with minor or no side effects, we evaluated intraurethral

Download English Version:

<https://daneshyari.com/en/article/9954768>

Download Persian Version:

<https://daneshyari.com/article/9954768>

[Daneshyari.com](https://daneshyari.com)